



Your Name / Adınız - Soyadınız

Your Signature / İmza

Student ID # / Öğrenci No

Professor's Name / Öğretim Üyesi

Your Department / Bölüm

- Calculators, cell phones off and away!.
- In order to receive credit, you must **show all of your work**. If you do not indicate the way in which you solved a problem, you may get little or no credit for it, even if your answer is correct. **Show your work in evaluating any limits, derivatives.**
- Place a box around your answer to each question.
- Use a **BLUE ball-point pen** to fill the cover sheet. Please make sure that your exam is complete.
- Time limit is 80 min.

Do not write in the table to the right.

Problem	Points	Score
1	25	
2	22	
3	23	
4	30	
Total:	100	

1. (a) 12 Points Find the point in which the line through $P(3, 2, 1)$ normal to the plane $2x - y + 2z = -2$ meets the plane.

Solution: Denote by \mathcal{L} the line through $P(3, 2, 1)$ normal to the plane $2x - y + 2z = -2$. Then write the parametric equations for \mathcal{L} .

$$\mathcal{L} : \begin{cases} x = 3 + 2t \\ y = 2 - t \\ z = 1 + 2t \end{cases}$$

So we are asked to find the point in which \mathcal{L} meets $2x - y + 2z = -2$. To this end, we simply substitute.


$$2(3 + 2t) - (2 - t) + 2(1 + 2t) = -2 \Rightarrow 6 + 6t - 2 + t + 2 + 4t = -2 \Rightarrow 9t = -8$$

So $t = -8/9$. Therefore the point where \mathcal{L} meets the plane $2x - y + 2z = -2$ has coordinates:

$$\mathcal{L} : \begin{cases} x = 3 + 2(-8/9) = 11/9 \\ y = 2 - (-8/9) = 26/9 \\ z = 1 + 2(-8/9) = -7/9 \end{cases}$$

That is, the point of intersection is $(11/9, 26/9, -7/9)$.

p.695, pr.37



- (b) 13 Points Find the volume of the parallelepiped (box) determined by the vectors $\mathbf{u} = \mathbf{i} + \mathbf{j} - 2\mathbf{k}$, $\mathbf{v} = -\mathbf{i} - \mathbf{k}$, $\mathbf{w} = 2\mathbf{i} + 4\mathbf{j} - 2\mathbf{k}$ when they are placed with the same initial point.


Solution: If $\mathbf{u} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$, $\mathbf{v} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$, $\mathbf{w} = c_1\mathbf{i} + c_2\mathbf{j} + c_3\mathbf{k}$, then

$$(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w} = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = (\mathbf{v} \times \mathbf{w}) \cdot \mathbf{u} = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix} = (\mathbf{w} \times \mathbf{u}) \cdot \mathbf{v}$$

which all have the same absolute value, since the interchanging two rows in a determinant does not change its absolute value, the volume of this parallelepiped is

$$|(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}| = \begin{vmatrix} 1 & 1 & -2 \\ -1 & 0 & -1 \\ 2 & 4 & -2 \end{vmatrix} = \begin{vmatrix} 0 & -1 \\ 4 & -2 \end{vmatrix} - \begin{vmatrix} -1 & -1 \\ 2 & -2 \end{vmatrix} + (-2) \begin{vmatrix} -1 & 0 \\ 2 & 4 \end{vmatrix} = 8$$

p.695, pr.37



2. (a) **10 Points** Evaluate the integral $\int e^\theta \sin \theta \, d\theta$.

Solution: We integrate by parts with a twist. Let $u = e^\theta$ and so $dv = \sin \theta \, d\theta$. Then $du = e^\theta \, d\theta$ and choose $v = -\cos \theta$. Therefore

$$\int e^\theta \sin \theta \, d\theta = \int u \, dv = uv - \int v \, du \quad (1)$$

$$= e^\theta (-\cos \theta) - \int (-\cos \theta) e^\theta \, d\theta \quad (2)$$

$$= -e^\theta \cos \theta + \int e^\theta \cos \theta \, d\theta \quad (3)$$

We next apply integration by parts to the integral on the right side of line (3). Letting

$$\int e^\theta \cos \theta \, d\theta = \left[\begin{array}{ll} u = e^\theta & dv = \cos \theta \, d\theta \\ du = e^\theta \, d\theta & v = \sin \theta \end{array} \right] = \underbrace{e^\theta}_{u} \underbrace{\sin \theta}_{v} - \int \underbrace{\sin \theta}_{v} \underbrace{e^\theta \, d\theta}_{du}$$

So we have

$$\int e^\theta \sin \theta \, d\theta = -e^\theta \cos \theta + e^\theta \sin \theta - \int e^\theta \sin \theta \, d\theta$$

Adding $\int e^\theta \sin \theta \, d\theta$ to both sides gives us

$$2 \int e^\theta \sin \theta \, d\theta = e^\theta (\sin \theta + \cos \theta).$$

Finally, dividing both sides by 2 and adding a constant of integration, we have

$$\int e^\theta \sin \theta \, d\theta = \boxed{\frac{1}{2} e^\theta (\sin \theta + \cos \theta) + c}$$

Incidentally, this integral can be evaluated by using $dv = e^\theta \, d\theta$ for both the first and the second applications of the integration by parts formula.

p.652, pr.3

- (b) **12 Points** Investigate the convergence/divergence of the series $\sum_{n=1}^{\infty} \frac{n}{n^2 + 1}$. Name the test you use.

○ Converges.

○ Diverges.

Test Used: _____

Solution: Use the Limit Comparison Test. Let $a_n = \frac{n}{n^2 + 1} > 0$ and choose $b_n = \frac{1}{n} > 0$. Then

$$\frac{a_n}{b_n} = \frac{\frac{n}{n^2 + 1}}{\frac{1}{n}} = \frac{n^2}{n^2 + 1} = \frac{n^2}{n^2 + 1} \cdot \frac{1/n^2}{1/n^2} = \frac{1}{1 + 1/n^2} \rightarrow \frac{1}{1 + 0} = 1$$

So $0 < L = 1 < \infty$ and this series *diverges*, because $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges.

We can also use Direct Comparison Test. Notice that if $n \geq 1$, then $n^2 + 1 \leq n^2 + n^2 = 2n^2$ and so

$$\frac{1}{n^2 + 1} \geq \frac{1}{2n^2} > 0 \Rightarrow a_n = \frac{1}{n^2 + 1} \geq \frac{1}{2n^2} = b_n > 0, \quad \forall n \geq 1.$$

Since $\sum_{n=1}^{\infty} \frac{1}{n}$ is harmonic series so it diverges. Therefore $\sum_{n=1}^{\infty} \frac{1}{2n}$ diverges too, as it is a scalar multiple of the harmonic series.

Hence the given series is a larger series than the series of scalar multiple series. We conclude by Direct Comparison Test that the given series diverges.

A third way could be the Integral Test. Define $f(x) = \frac{1}{x^2 + 1}$ is positive, continuous and decreasing for all $x \geq 1$. So we compute the improper integral

$$\int_1^{\infty} \frac{x}{x^2 + 1} dx = \lim_{b \rightarrow \infty} \int_1^b \frac{x}{x^2 + 1} dx = \lim_{b \rightarrow \infty} \left[\frac{1}{2} \ln(x^2 + 1) \right]_1^b = \lim_{b \rightarrow \infty} \left[\frac{1}{2} \ln(b^2 + 1) - \frac{1}{2} \ln 2 \right]$$

Therefore the integral $\int_1^{\infty} \frac{x}{x^2 + 1} dx$ diverges. By the Integral Test, the series diverges too.

p.533, pr.95

3. (a) **13 Points** Find the values of $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ at $(1, 1, 1)$ if

$$z^3 - xy + yz + y^3 - 2 = 0.$$

Solution: We can do this in two ways.

First we can differentiate implicitly with respect to x (y is held constant).

$$\begin{aligned} \frac{\partial}{\partial x} (z^3 - xy + yz + y^3 - 2) &= \frac{\partial}{\partial x} (0) \\ 3z^2 \frac{\partial z}{\partial x} - y + y \frac{\partial z}{\partial x} &= 0 \\ (3z^2 + y) \frac{\partial z}{\partial x} &= y \\ \frac{\partial z}{\partial x} &= \frac{y}{3z^2 + y} \\ \Rightarrow \frac{\partial z}{\partial x} \Big|_{(1,1,1)} &= \frac{1}{3(1)^2 + 1} = \frac{1}{4}. \end{aligned}$$

Similarly, differentiating with respect to y (treating x constant), we have

$$\begin{aligned} \frac{\partial}{\partial y} (z^3 - xy + yz + y^3 - 2) &= \frac{\partial}{\partial y} (0) \\ 3z^2 \frac{\partial z}{\partial y} - x + z + y \frac{\partial z}{\partial y} + 3y^2 &= 0 \\ (3z^2 + y) \frac{\partial z}{\partial y} &= x - z - 3y^2 \\ \frac{\partial z}{\partial y} &= \frac{x - z - 3y^2}{3z^2 + y} \\ \Rightarrow \frac{\partial z}{\partial y} \Big|_{(1,1,1)} &= \frac{1 - 1 - 3(1)^2}{3(1)^2 + 1} = \frac{-3}{4}. \end{aligned}$$

As an alternative method, let $F(x, y, z) = z^3 - xy + yz + y^3 - 2 = 0$. Then $F_x(x, y, z) = -y$, $F_y(x, y, z) = -x + z + 3y^2$ and $F_z(x, y, z) = 3z^2 + y$. Therefore, by implicit differentiation formulas (Theorem 8, page 780 of the textbook)

$$\begin{aligned} \frac{\partial z}{\partial x} \Big|_{(1,1,1)} &= -\frac{F_x}{F_z} \Big|_{(1,1,1)} = -\frac{-y}{3z^2 + y} \Big|_{(1,1,1)} = \frac{1}{4} \\ \frac{\partial z}{\partial y} \Big|_{(1,1,1)} &= -\frac{F_y}{F_z} \Big|_{(1,1,1)} = -\frac{-x + z + 3y^2}{3z^2 + y} \Big|_{(1,1,1)} = \frac{-3}{4}. \end{aligned}$$

p.879, pr.42

- (b) **10 Points** Let $z = \sqrt{y - x}$. Find the equation for tangent plane to this surface at the point $(1, 2, 1)$.

Solution: Let $F(x, y, z) = \sqrt{y - x} - z$. Then

$$\begin{aligned} F_x(x, y, z) &= -\frac{1}{2}(y - x)^{-1/2} \\ F_y(x, y, z) &= \frac{1}{2}(y - x)^{-1/2} \\ F_z(x, y, z) &= -1 \end{aligned}$$

At $(1, 2, 1)$, we have

$$F_x(1, 2, 1) = -\frac{1}{2}(2-1)^{-1/2} = -\frac{1}{2}$$

$$F_y(1, 2, 1) = \frac{1}{2}(2-1)^{-1/2} = \frac{1}{2}$$

$$F_z(1, 2, 1) = -1.$$

Hence the tangent plane at $(1, 2, 1)$ is

$$-\frac{1}{2}(x-1) + \frac{1}{2}(y-2) - (z-1) = 0 \Rightarrow \boxed{x - y + 2z = 1}$$

p.879, pr.34

4. (a) **15 Points** Find two numbers a and b with $a \leq b$ such that $\int_a^b (6-x-x^2) dx$ has its largest value.

Solution: Let $F(a, b) = \int_a^b (6-x-x^2) dx$ where $a \leq b$. the boundary of the domain of F is the $a = b$ in the ab -plane, and $F(a, a) = 0$, so F is identically 0 on the boundary of its domain. For interior critical points we have:

$$\frac{\partial F}{\partial a} = -(6-a-a^2) = 0 \Rightarrow -(3+a)(2-a) = 0 \Rightarrow a = -3, 2$$

$$\frac{\partial F}{\partial b} = -(6-b-b^2) = 0 \Rightarrow -(3+b)(2-b) = 0 \Rightarrow b = -3, 2$$

Hence the candidates for critical points are $(-3, -3)$, $(-3, 2)$, $(2, -3)$, and $(2, 2)$. Since $a \leq b$, only one of these four points is a critical point, namely, it is $(-3, 2)$. Next

$$F(-3, 2) = \int_{-3}^2 (6-x-x^2) dx = \left[6x - \frac{x^2}{2} - \frac{x^3}{3} \right]_{-3}^2 = 12 - 2 - \frac{8}{3} - \left(-18 - \frac{9}{2} + 9 \right) = 75/3$$

is the maximum value of F and gives the area under the parabola $y = 6-x-x^2$ that is above the x -axis. Therefore $a = -3$ and $b = 2$.

p.695, pr.37

- (b) **15 Points** Find the absolute maximum and minimum values of $f(x, y) = x^2 + xy + y^2 - 3x + 3y$ on the given triangular region R .

Solution:

• On AB , we have $f(x, y) = f(x, 4-x) = x^2 - 10x + 28$ for $0 \leq x \leq 4$. So $f'(x) = 2x - 10 \Rightarrow x = 5 \notin [0, 4] \Rightarrow$ no critical points in the interior of AB .

Endpoints of AB : $f(4, 0) = 4$ and $f(0, 4) = 28$.

• On OB , we have $f(x, y) = f(x, 0) = x^2 - 3x$ for $0 \leq x \leq 4$. So $f'(x) = 2x - 3 = 0 \Rightarrow x = 3/2$ and $y = 0 \Rightarrow (3/2, 0)$ is an interior critical point of OB with $f(3/2, 0) = -9/4$.

Endpoints of BC : $f(0, 0) = 0$ and $f(4, 0) = 4$.

• On OA , we have $f(x, y) = f(0, y) = y^2 + 3y$ for $0 \leq y \leq 4$. So $f'(y) = 2y + 3 = 0 \Rightarrow y = -3/2$ and $x = 0 \Rightarrow (0, -3/2)$ is not on the segment OA . So no interior point occurs as C.P.

Endpoints of OA : $f(0, 0) = 0$ and $f(0, 4) = 28$.

• Interior Points of this triangular region R : $f_x(x, y) = 2x + y - 3 = 0$ and $f_y(x, y) = x + 2y + 3 = 0 \Rightarrow x = 3, y = -3 \Rightarrow (3, -3)$. But $(3, -3)$ is not an interior of R . So f has no interior critical point.

Therefore *abs. max* is 28 at $(0, 4)$ and the *abs. min.* is $-9/4$ at $(3/2, 0)$.

p.695, pr.37

