



Your Name / Adınız - Soyadınız

Your Signature / İmza

Student ID # / Öğrenci No

--	--	--	--	--	--	--	--	--	--

Professor's Name / Öğretim Üyesi

Your Department / Bölüm

- This exam is closed book.
- Give your answers in exact form (for example $\frac{\pi}{3}$ or $5\sqrt{3}$), except as noted in particular problems.
- Calculators, cell phones are not allowed.
- In order to receive credit, you must **show all of your work**. If you do not indicate the way in which you solved a problem, you may get little or no credit for it, even if your answer is correct. **Show your work in evaluating any limits, derivatives.**
- Place a box around your answer to each question.
- If you need more room, use the backs of the pages and indicate that you have done so.
- Do not ask invigilator anything.
- This exam has 5 pages plus this cover sheet and 6 problems. Please make sure that your exam is complete.

Problem	Points	Score
1	20	
2	15	
3	15	
4	15	
5	15	
6	20	
Total:	100	

. Time limit is 90 min.

Do not write in the table to the right.

1. Suppose $f(x) = \frac{x^2}{x^2-1}$, $f'(x) = \frac{-2x}{(x^2-1)^2}$, $f''(x) = \frac{6x^2+2}{(x^2-1)^3}$, p.212, pr.80

- (a) **4 Points** All critical points of f , and the intervals where f is increasing and decreasing;

Solution: The critical points are only when $f'(x) = \frac{-2x}{(x^2-1)^2} = 0$, that is when $2x = 0$ and so the point is at $x = 0$. This splits the real line into 4 open subintervals, namely, $(-\infty, -1)$, $(-1, 0)$, $(0, 1)$, and $(1, \infty)$. By considering test values on each of these intervals, we see that f is increasing on $(-\infty, -1) \cup (-1, 0)$ and decreasing on $(0, 1) \cup (1, \infty)$.

- (b) **4 Points** All inflection points of f , and the open intervals where f is concave up resp. concave down.

Solution: Since $f''(x) = \frac{6x^2+2}{(x^2-1)^3} \neq 0$ and there is no domain point for f where the second derivative is undefined, there are no points of inflection for the graph of f . By looking at the sign for f'' , we see that the graph is concave up on $(-\infty, -1) \cup (1, \infty)$ and concave down on $(-1, 1)$.

- (c) **4 Points** Classify the critical points of f as either local maxima, local minima or neither.


Solution: At $x = 0$, the graph has a local maximum, namely, the point $(0, 0)$ is a point of local maximum and there is no point of local minimum.

- (d) **4 Points** All asymptotes of f .

Solution: There are three asymptotes. First $x = -1$ and $x = 1$ are vertical asymptotes since $\lim_{x \rightarrow \pm 1^\pm} f(x) = +\infty$. Next $y = 1$ is the horizontal asymptote, since $\frac{x^2}{x^2-1} \rightarrow 1$ as $x \rightarrow \pm\infty$.

- (e) **4 Points** Sketch the graph of f using your results in (a), (b), (c) and (d).


Solution:



2. The rectangle shown here has one side on the positive y -axis, one side on the positive x -axis, and its upper right-hand vertex on the curve $y = e^{-x^2}$. p.432, pr.121

- (a) **5 Points** Write the area of the rectangle in terms of a .

Solution: The area is $A = ab = ae^{-a^2}$.



- (b) **10 Points** What dimensions give the rectangle its largest area, and what is that area?

Solution: From part (a), we have $A = ae^{-a^2}$. Differentiating with respect to a gives

$$\frac{dA}{da} = e^{-a^2} + (a)(-2a)e^{-a^2} = e^{-a^2}(1 - 2a^2).$$


Hence $\frac{dA}{da} < 0$ for $a > \frac{1}{\sqrt{2}}$ and $\frac{dA}{da} > 0$ for $0 < a < \frac{1}{\sqrt{2}}$.

Solving $\frac{dA}{da} = 0$ for a , we have absolute maximum of $\frac{1}{\sqrt{2}}e^{-1/2} = \frac{1}{\sqrt{2e}}$ at $a = \frac{1}{\sqrt{2}}$ units long by $b = e^{-1/2} = \frac{1}{\sqrt{e}}$ units high.

3. **15 Points** Use the method of disks to find the volume of the solid generated by revolving, about the line x -axis, the region bounded by: $y = \frac{1}{\sqrt{1+x^2}}$, $y = 0$, $\frac{-1}{\sqrt{3}} \leq x \leq \sqrt{3}$. p.415, pr.121

Solution:

$$\begin{aligned} V &= \pi \int_{-1/\sqrt{3}}^{\sqrt{3}} \left(\frac{1}{\sqrt{1+x^2}} \right)^2 dx \\ &= \pi \int_{-1/\sqrt{3}}^{\sqrt{3}} \frac{1}{1+x^2} dx = \pi [\tan^{-1} x]_{-1/\sqrt{3}}^{\sqrt{3}} \\ &= \pi \left[\tan^{-1} \sqrt{3} - \tan^{-1} \left(-\frac{\sqrt{3}}{3} \right) \right] \\ &= \pi \left[\frac{\pi}{3} - \left(-\frac{\pi}{6} \right) \right] = \frac{\pi^2}{2} \end{aligned}$$



4. 15 Points Find the length of the curve given by $x = y^{2/3}$, $1 \leq y \leq 8$ p.298 pr.39

Solution:

$$\begin{aligned}x &= y^{2/3} \Rightarrow \frac{dx}{dy} = \frac{2}{3}y^{-1/3} \Rightarrow \left(\frac{dx}{dy}\right)^2 = \frac{4y^{-2/3}}{9} \Rightarrow L = \int_1^8 \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy \\L &= \int_1^8 \sqrt{1 + \frac{4}{9y^{2/3}}} dy = \int_1^8 \frac{\sqrt{9y^{2/3} + 4}}{3y^{1/3}} dy = \frac{1}{3} \int_1^8 \sqrt{9y^{2/3} + 4} (y^{-1/3}) dy \\; [u &= 9y^{2/3} + 4 \Rightarrow du = 6y^{-1/3} dy; y = 1 \Rightarrow u = 13, y = 8 \Rightarrow u = 40] \\ \rightarrow L &= \frac{1}{18} \int_{13}^{40} u^{1/2} du = \frac{1}{18} \left[\frac{2}{3} u^{3/2} \right]_{13}^{40} = \frac{1}{27} [40^{3/2} - 13^{3/2}] \approx 7.634\end{aligned}$$

5. (a) 6 Points If $9e^{2y} = x^2$, then solve for y in terms of x . p.431, pr.81

Solution: We isolate y .

$$\begin{aligned} 9e^{2y} &= x^2 \Rightarrow e^{2y} = \frac{x^2}{9} \Rightarrow \ln e^{2y} = \ln \frac{x^2}{9} \Rightarrow 2y \ln e = \ln \frac{x^2}{9} \\ &\Rightarrow y = \frac{1}{2} \ln \frac{x^2}{9} = \ln \sqrt{\frac{x^2}{9}} = \ln \left| \frac{x}{3} \right| \\ y &= \ln |x| - \ln 3 \end{aligned}$$

- (b) 9 Points $\lim_{x \rightarrow 0} \frac{5 - 5 \cos x}{e^x - x - 1} = ?$ p.431, pr.101

Solution: This has the indeterminate form $\frac{0}{0}$. Hence the L'Hôpital's Rule applies.

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{5 - 5 \cos x}{e^x - x - 1} &\stackrel{\text{L'H}}{=} \lim_{x \rightarrow 0} \frac{5 \sin x}{e^x - 1} \stackrel{\text{L'H}}{=} \lim_{x \rightarrow 0} \frac{5 \cos x}{e^x} \\ &= \frac{5 \cos(0)}{e^0} = 5 \end{aligned}$$

6. Evaluate the following integrals.

- (a) 10 Points $\int_{-\ln 4}^{-\ln 2} 2e^\theta \cosh \theta \, d\theta$ p.431, pr.53

Solution: Since $\cosh \theta = \frac{e^\theta + e^{-\theta}}{2}$, we have

$$\begin{aligned} \int_{-\ln 4}^{-\ln 2} 2e^\theta \cosh \theta \, d\theta &= \int_{-\ln 4}^{-\ln 2} 2e^\theta \left(\frac{e^\theta + e^{-\theta}}{2} \right) d\theta = \int_{-\ln 4}^{-\ln 2} (e^{2\theta} + 1) d\theta = \left[\frac{e^{2\theta}}{2} + \theta \right]_{-\ln 4}^{-\ln 2} \\ &= \left(\frac{e^{-2 \ln 2}}{2} - \ln 2 \right) - \left(\frac{e^{-2 \ln 4}}{2} - \ln 4 \right) = \left(\frac{1}{8} - \ln 2 \right) - \left(\frac{1}{32} - \ln 4 \right) \\ &= \frac{3}{32} - \ln 2 + 2 \ln 2 = \frac{3}{32} + \ln 2 \end{aligned}$$

- (b) 10 Points $\int \frac{dx}{\sqrt{9-4x^2}}$ p.431, pr.65

Solution: We use the substitution $u = \frac{2x}{3}$ and so $du = \frac{2}{3}dx$.

$$\begin{aligned}\int \frac{dx}{\sqrt{9-4x^2}} &= \frac{1}{3} \int \frac{dx}{\sqrt{1-\left(\frac{2x}{3}\right)^2}} \\ &= \frac{1}{2} \int \frac{du}{\sqrt{1-u^2}}, \text{ where } u = \frac{2x}{3}, du = \frac{2}{3}dx; \\ &= \frac{1}{2} \sin^{-1}\left(\frac{2x}{3}\right) + C\end{aligned}$$

