

Your Name

Your Signature

Student ID #

--	--	--	--	--	--	--	--	--

Professor's Name

Your Department

- This exam is closed book.
- Give your answers in exact form (for example $\frac{\pi}{3}$ or $5\sqrt{3}$), except as noted in particular problems.
- Calculators, cell phones are not allowed.
- In order to receive credit, you must **show all of your work**. If you do not indicate the way in which you solved a problem, you may get little or no credit for it, even if your answer is correct. **Show your work in evaluating any limits, derivatives.**
- Place to each question.
- If you need more room, use the backs of the pages and indicate that you have done so.
- Raise your hand if you have a question.
- This exam has ?? pages, plus this cover sheet. Please make sure that your exam is complete.

Question	Points	Score
1	??	
2	??	
3	??	
4	??	

Question	Points	Score
5	??	
6	??	
7	??	
Total	??	

1. (?? total points) Given the function $f(x) = \frac{x^2 + 6x + 5}{x + 5}$, and point $x_0 = -5$ and $\varepsilon = 0.05$ p.64, pr.34

(a) (5 Points) Find $L = \lim_{x \rightarrow -5} f(x)$. (DO NOT USE L'HÔPITAL'S RULE)

$$\begin{aligned} \lim_{x \rightarrow -5} \frac{x^2 + 6x + 5}{x + 5} &= \lim_{x \rightarrow -5} \frac{(x+5)(x+1)}{(x+5)} \\ &= \lim_{x \rightarrow -5} (x+1) \\ &= -4, \quad x \neq -5. \end{aligned}$$

(b) (10 Points) Find a number δ such that for all x

$$0 < |x - x_0| < \delta \Rightarrow |f(x) - L| < \varepsilon.$$

Step 1

$$\begin{aligned} \left| \left(\frac{x^2 + 6x + 5}{x + 5} \right) - (-4) \right| < 0.05 &\Rightarrow -0.05 < \frac{x^2 + 6x + 5}{x + 5} + 4 < 0.05 \\ &\Rightarrow -4.05 < x + 1 < -3.95, \quad x \neq -5 \\ &\Rightarrow -5.05 < x < -4.95, \quad x \neq -5 \end{aligned}$$

Step 2

$$\begin{aligned} |x - (-5)| < \delta &\Rightarrow -\delta < x + 5 < \delta \Rightarrow -\delta - 5 < x < \delta - 5. \\ \text{Then } -\delta - 5 = -5.05 &\Rightarrow \delta = 0.05, \text{ or } \delta - 5 = -4.95 \Rightarrow \delta = 0.05; \text{ thus } \delta = 0.05. \end{aligned}$$

2. Evaluate the following limits.

(a) (10 Points) $\lim_{x \rightarrow 1} \frac{x^{1/3} - 1}{\sqrt{x} - 1}$ (DO NOT USE L'HÔPITAL'S RULE)

$$\begin{aligned} \lim_{x \rightarrow 1} \frac{x^{1/3} - 1}{\sqrt{x} - 1} &= \lim_{x \rightarrow 1} \frac{x^{1/3} - 1}{\sqrt{x} - 1} \frac{(x^{2/3} + x^{1/3} + 1)(\sqrt{x} + 1)}{(x^{2/3} + x^{1/3} + 1)(\sqrt{x} + 1)} \\ &= \lim_{x \rightarrow 1} \frac{(x-1)(\sqrt{x} + 1)}{(x-1)(x^{2/3} + x^{1/3} + 1)} \\ &= \lim_{x \rightarrow 1} \frac{\sqrt{x} + 1}{x^{2/3} + x^{1/3} + 1} \\ &= \frac{1 + 1}{1 + 1 + 1} \\ &= \frac{2}{3} \end{aligned}$$

(b) (10 Points) $\lim_{x \rightarrow 0} \frac{x \sin x}{2 - 2 \cos x}$ p.178 pr.101 (DO NOT USE L'HÔPITAL'S RULE)

$$\begin{aligned} \lim_{x \rightarrow 0} \frac{x \sin x}{2 - 2 \cos x} &= \lim_{x \rightarrow 0} \frac{x \sin x}{2(1 - \cos x)} = \lim_{x \rightarrow 0} \frac{x \sin x}{2(2 \sin^2(\frac{x}{2}))} = \lim_{x \rightarrow 0} \left[\frac{\frac{x}{2} \cdot \frac{x}{2}}{\sin^2(\frac{x}{2})} \cdot \frac{\sin x}{x} \right] \\ &= \lim_{x \rightarrow 0} \left[\frac{\frac{x}{2}}{\sin(\frac{x}{2})} \cdot \frac{\frac{x}{2}}{\sin(\frac{x}{2})} \cdot \frac{\sin x}{x} \right] \\ &= (1)(1)(1) = 1. \end{aligned}$$

3. (10 Points) Find $\frac{dy}{dt}$ if $y = (t^{-3/4} \sin t)^{4/3}$.

p.147 pr.46

$$\begin{aligned}\frac{dy}{dt} &= \frac{d}{dt} \left(t^{-3/4} \sin t \right)^{4/3} \\ &= \frac{4}{3} \left(t^{-3/4} \sin t \right)^{1/3} \left[-\frac{3}{4} t^{-7/4} \sin t \right. \\ &\quad \left. + t^{-3/4} \cos t \right]\end{aligned}$$

4. (15 Points) Find the equations of normals to the curve

$$xy + 2x - y = 0$$

that are parallel to the line $2x + y = 0$.

p.154, pr.40

$$\begin{aligned}xy + 2x - y = 0 &\implies x \frac{dy}{dx} + y + 2 - \frac{dy}{dx} = 0 \\ &\implies \frac{dy}{dx} = \frac{y+2}{1-x};\end{aligned}$$

the slope of the line $2x + y = 0$ is -2 . In order to be parallel, the normal lines must also have slope of -2 . Since a normal is perpendicular to a tangent, the slope of tangent is $\frac{1}{2}$. Therefore

$$\begin{aligned}\frac{y+2}{1-x} &= \frac{1}{2} \implies 2y+4 = 1-x \\ &\implies x = -3 - 2y.\end{aligned}$$

Substituting in the original equation,

$$\begin{aligned}y(-3-2y) + 2(-3-2y) &= 0 \\ &\implies y^2 + 4y + 3 = 0 \\ &\implies y = -3 \text{ or } y = -1.\end{aligned}$$

$$\begin{aligned}\text{If } y &= -3, \text{ then } x = 3 \text{ and } y + 3 = -2(x - 3) \\ &\implies y = -2x + 3.\end{aligned}$$

$$\begin{aligned}\text{If } y &= -1, \text{ then } x = -1 \text{ and } y + 1 = -2(x + 1) \\ &\implies y = -2x - 3.\end{aligned}$$

5. (15 Points) For what values of a and b is

$$f(x) = \begin{cases} -2 & x \leq -1 \\ ax - b, & -1 < x < 1 \\ 3, & x \geq 1 \end{cases}$$

continuous at every x .

p.83, pr.45

Clearly f is continuous if $x \neq -1$ and for $x \neq 1$ for if $x < -1$ or if $-1 < x < 1$ or if $x > 1$, f is a polynomial, regardless the values of a and b . For continuity at $x = -1$, we require that the one-sided limits of $f(x)$ at $x = -1$ be equal. But $\lim_{x \rightarrow -1^-} f(x) = -2$ and $\lim_{x \rightarrow -1^+} f(x) = a(-1) + b = -a + b$.

Similarly, for continuity at $x = 1$, we require that the one-sided limits of $f(x)$ at $x = 1$ be equal. But $\lim_{x \rightarrow 1^-} f(x) = a(1) + b = a + b$ and $\lim_{x \rightarrow 1^+} f(x) = 3$.

Equality of one-sided limits is equivalent to

$$\begin{aligned}-2 &= -a + b \text{ and } a + b = 3 \\ &\implies a = \frac{5}{2} \text{ and } b = \frac{1}{2}.\end{aligned}$$

6. (15 Points) Assume that $f(x)$ and $g(x)$ are differentiable functions satisfying

$$\begin{aligned} g(0) &= 1 & f(0) &= 1 & f(1) &= 3 & g(1) &= 5 \\ g'(0) &= \frac{1}{2} & f'(0) &= -3 & f'(1) &= \frac{1}{2} & g'(1) &= -4 \end{aligned}$$

Let $h(x) = f(x + g(x))$. Evaluate $h'(0)$. p.148, pr.74

First, by the Chain Rule, we have $h'(x) = f'(x + g(x))(1 + g'(x))$. Then $h'(0) = f'(0 + g(0))(1 + g'(0)) = f'(g(0))(1 + g'(0))$.

$$\text{Hence } h'(0) = f'(1)\left(1 + \frac{1}{2}\right) = \left(\frac{1}{2}\right)\left(\frac{3}{2}\right) = \frac{3}{4}.$$

7. (10 Points) Find the value or values of c that satisfy the equation

$$\frac{f(b) - f(a)}{b - a} = f'(c)$$

in the conclusion of the Mean Value Theorem for the function $f(x) = x + \frac{1}{x}$ and interval $[\frac{1}{2}, 2]$. p.196, pr.3

When $f(x) = x + \frac{1}{x}$ for $\frac{1}{2} \leq x \leq 2$, then

$$\frac{f(2) - f(1/2)}{2 - 1/2} = f'(c) \Rightarrow 0 = 1 - \frac{1}{c^2} \Rightarrow c = \pm 1$$

But $-1 \notin [\frac{1}{2}, 2]$, so $c = 1$.