

Your Name / Ad - Soyad

(90 min.)

Signature / İmza

Student ID # / Öğrenci No

(mavi tükenmez!)

Problem	1	2	3	4	Total
Points:	40	25	26	24	115
Score:					

You have **90 minutes**. (Cell phones off and away!). No books, notes or calculators are permitted. **Mysterious or unsupported answers will not receive full credit.** A correct answer, unsupported by calculations, explanation, or algebraic work will receive no credit; an incorrect answer supported by substantially correct calculations and explanations might still receive partial credit.

1. (a) (10 Points) Evaluate the integral $\int x^3 e^{(x^2)} dx$.

Solution: Let $y = x^2$. Then $dy = 2x dx$. Therefore

$$\begin{aligned}\int x^3 e^{(x^2)} dx &= \frac{1}{2} \int x^2 e^{(x^2)} 2x dx \\ &= \frac{1}{2} \int y e^y dy\end{aligned}$$

We now integrate by parts. For let $u = y$ and so $dv = e^y dy$. Then $du = dy$ and choose $v = e^y$. Hence

$$\begin{aligned}\int y e^y dy &= \int u dv = uv - \int v du \\ &= y e^y - \int e^y dy \\ &= y e^y - e^y + C_1\end{aligned}$$

Finally, we have

$$\int x^3 e^{(x^2)} dx = \frac{1}{2} (x^2 e^{x^2} - e^{x^2} + C_1)$$

p.491, pr.86

- (b) (10 Points) Find the parametric equations of the line normal to the surface $x^2 z + x z^2 + y^2 = yz + 5x + 5$ at the point $P_0(1, 2, 3)$.

Solution: Let $F(x, y, z) = x^2 z + x z^2 + y^2 - yz - 5x - 5 = 0$. Then

$$\begin{aligned}F_x &= 2xz + z^2 - 5 \\ F_y &= 2y - z \\ F_z &= x^2 + 2xz - y.\end{aligned}$$

Then

$$\begin{aligned}F_x(1, 2, 3) &= 6 + 9 - 5 = 10 \\ F_y(1, 2, 3) &= 4 - 3 = 1 \\ F_z(1, 2, 3) &= 1 + 6 - 2 = 5.\end{aligned}$$

The parametric equations for required normal line are

$$x = 1 + 8t, \quad y = 2 + t, \quad z = 3 + 5t$$

p.491, pr.86

- (c) (10 Points) Evaluate the integral $\int \frac{\ln x}{x + x \ln x} dx$.

Solution: Let $y = \ln x$ and so $dy = \frac{1}{x} dx$. Then

$$\begin{aligned}\int \frac{\ln x}{x + x \ln x} dx &= \int \frac{\ln x}{1 + \ln x} \frac{1}{x} dx \\ &= \int \frac{y}{1 + y} dy \\ &= \int \frac{y + 1 - 1}{1 + y} dy \\ &= \int \left(1 - \frac{1}{1 + y} \right) dy \\ &= y - \ln|1 + y| + C = \ln x - \ln|1 + \ln x| + C\end{aligned}$$

p.491, pr.107

- (d) (10 Points) Use logarithmic differentiation to find the derivative of

$$y = \sqrt[3]{\frac{x(x+1)(x+2)}{(x^2+1)(2x+3)}}$$

with respect to x .

Solution:

$$\begin{aligned}\ln y &= \frac{1}{3} [\ln x + \ln(x+1) + \ln(x+2) - \ln(x^2+1) - \ln(2x+3)] \\ \frac{1}{y} \frac{dy}{dx} &= \frac{1}{3} \left[\frac{1}{x} + \frac{1}{x+1} + \frac{1}{x+2} - \frac{2x}{x^2+1} - \frac{2}{2x+3} \right] \\ \frac{dy}{dx} &= \frac{y}{3} \left[\frac{1}{x} + \frac{1}{x+1} + \frac{1}{x+2} - \frac{2x}{x^2+1} - \frac{2}{2x+3} \right] \\ &= \frac{1}{3} \sqrt[3]{\frac{x(x+1)(x+2)}{(x^2+1)(2x+3)}} \left[\frac{1}{x} + \frac{1}{x+1} + \frac{1}{x+2} - \frac{2x}{x^2+1} - \frac{2}{2x+3} \right]\end{aligned}$$

p.94, pr.34

2. Given the three points $P(1,0,1)$, $Q(2,0,0)$, and $R(-1,2,2)$.

- (a) (6 Points) Find the area of the triangle having P , Q and R as vertices.

Solution: First form the vectors $\vec{PQ} = (2-1)\mathbf{i} + (0-0)\mathbf{j} + (0-1)\mathbf{k} = \mathbf{i} - \mathbf{k}$ and $\vec{PR} = (-1-1)\mathbf{i} + (2-0)\mathbf{j} + (2-1)\mathbf{k} = -2\mathbf{i} + 2\mathbf{j} + \mathbf{k}$

$$A = \frac{1}{2} \|\vec{PQ} \times \vec{PR}\|$$

$$\vec{PQ} \times \vec{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & -1 \\ -2 & 2 & 1 \end{vmatrix}$$

$$= 2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

$$\|\vec{PQ} \times \vec{PR}\| = \sqrt{2^2 + 1^2 + 2^2} = \sqrt{9} = 3$$

$$A = \frac{3}{2}$$

p.551, pr.32

- (b) (6 Points) Find the angle (in degrees) at the vertex P of the triangle having P , Q and R as vertices.

Solution: Let θ denote the angle we want to find. Then

$$\cos \theta = \frac{\vec{PQ} \cdot \vec{PR}}{\|\vec{PQ}\| \|\vec{PR}\|}$$

$$= \frac{(\mathbf{i} - \mathbf{k}) \cdot (-2\mathbf{i} + 2\mathbf{j} + \mathbf{k})}{\sqrt{2}\sqrt{9}}$$

$$= \frac{1}{3\sqrt{2}} ((1)(-2) + (0)(2) + (-1)(1))$$

$$= \frac{-3}{3\sqrt{2}} = -\frac{1}{\sqrt{2}}$$

Hence $\theta = 135$ degrees.

p.551, pr.32

- (c) (6 Points) Find the equation of the plane containing the points P , Q and R .

Solution: We know from part (a) that the vector

$$\mathbf{n} = 2\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

is normal to the plane. Therefore

$$\mathbf{n} \cdot \vec{P_0P} =$$

gives

$$(2\mathbf{i} + \mathbf{j} + 2\mathbf{k}) \cdot ((x-1)\mathbf{i} + (y-0)\mathbf{j} + (z-1)\mathbf{k}) = 0$$

$$\text{Hence } 2(x-1) + y + 2(z-1) = 0 \Rightarrow 2x + y + 2z = 4$$

p.551, pr.32

- (d) (7 Points) Find the value(s) of c if the function

$$f(x,y) = \begin{cases} \frac{3x^2y}{x^2+y^2} & (x,y) \neq (0,0) \\ c, & (x,y) = (0,0) \end{cases}$$

is continuous at $(0,0)$.

Solution: We employ the polar coordinates: $x = r \cos \theta$ and $y = r \sin \theta$. Then $x^2 + y^2 = r^2$ and $r \rightarrow 0$ as $(x,y) \rightarrow (0,0)$. Hence we have

$$\lim_{(x,y) \rightarrow (0,0)} f(x,y) = \lim_{r \rightarrow 0} \frac{3r^3 \cos^2 \theta \sin \theta}{r^2}$$

$$= \lim_{r \rightarrow 0} (3r \cos^2 \theta \sin \theta)$$

$$= 0$$

So $f(x,y)$ is continuous at $(0,0)$ iff $c=0$.

p.82, pr.35

3. (a) (10 Points) Find $\frac{\partial z}{\partial u}$ and $\frac{\partial z}{\partial v}$ when $u = \ln 2$, $v = 1$ if $z = 5 \tan^{-1} x$ and $x = e^u + \ln v$.

Solution: We apply the two chain rule formulas.

$$\frac{\partial z}{\partial u} = \frac{dz}{dx} \frac{\partial x}{\partial u} \quad \text{and} \quad \frac{\partial z}{\partial v} = \frac{dz}{dx} \frac{\partial x}{\partial v}.$$

Differentiating gives

$$\frac{dz}{dx} = \frac{5}{1+x^2}, \quad \frac{\partial x}{\partial u} = e^u, \quad \frac{\partial x}{\partial v} = \frac{1}{v}.$$

Moreover, when $u = \ln 2$ and $v = 1$, we have $x = e^{\ln 2} + \ln 1 = 2 + 0 = 2$. And derivatives at these points have values:

$$\left. \frac{dz}{dx} \right|_{x=2} = \frac{5}{1+2^2} = 1, \quad \left. \frac{\partial x}{\partial u} \right|_{u=\ln 2} = e^{\ln 2} = 2, \quad \left. \frac{\partial x}{\partial v} \right|_{v=1} = \frac{1}{1} = 1.$$

Therefore

$$\boxed{\left. \frac{\partial z}{\partial u} \right|_{u=\ln 2, v=1} = (1)(2) = 2}, \quad \boxed{\left. \frac{\partial z}{\partial v} \right|_{u=\ln 2, v=1} = (1)(1) = 1}.$$

p.72, pr.8

- (b) (16 Points) Find the absolute maximum and minimum values of $f(x, y) = x^2 - y^2 - 2x + 4y$ on the given region R .

Solution:

- *Interior Points of this triangular region R :* $f_x(x, y) = 2x - 2 = 0 \Rightarrow x = 1$ and $f_y(x, y) = -2y + 4 = 0 \Rightarrow y = 2 \Rightarrow (1, 2)$ is an interior critical point of R with $f(1, 2) = 3$.


- *On AB ,* we have $f(x, x+2) = -2x+4$ for $-2 \leq x \leq 2$. So $f'(x) = -2 = 0 \Rightarrow$ no critical points in the interior of AB . Endpoints of AB : $f(-2, 0) = 8$ and $f(2, 4) = 0$.

- *On BC ,* we have $f(x, y) = f(2, y) = -y^2 + 4y$ for $0 \leq y \leq 4$. So $f'(2, y) = -2y + 4 = 0 \Rightarrow y = 2$ and $x = 2 \Rightarrow (2, 2)$ is an interior critical point of BC with $f(2, 2) = 4$. Endpoints of BC : $f(2, 0) = 0$ and $f(2, 4) = 0$.

- *On AC ,* we have $f(x, y) = f(x, 0) = x^2 - 2x$ for $-2 \leq x \leq 2$. So $f'(x, 0) = 2x - 2 = 0 \Rightarrow x = 1$ and $y = 0 \Rightarrow (1, 0)$ is an interior critical point of AC with $f(1, 0) = -1$. Endpoints of AC : $f(-2, 0) = 8$ and $f(2, 0) = 0$.

- Therefore the absolute maximum is 8 at $(-2, 0)$ and the absolute minimum is -1 at $(1, 0)$,

p.317, pr.33



4. (a) (10 Points) Use the fact that $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$ to find the *first three nonzero terms* of the Maclaurin series for $\int \frac{e^x - 1}{x} dx$.

Solution: Using

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \cdots, \quad -\infty < x < \infty$$

we have

$$\frac{e^x - 1}{x} = 1 + \frac{x}{2!} + \frac{x^2}{3!} + \frac{x^3}{4!} + \frac{x^4}{5!} + \cdots, \quad -\infty < x < \infty$$

This is the Maclaurin series for $\frac{e^x - 1}{x}$. We can now do term-by-term integration.

$$\int \left(\frac{e^x - 1}{x} \right) dx = \int \left(1 + \frac{x}{2!} + \frac{x^2}{3!} + \frac{x^3}{4!} + \frac{x^4}{5!} + \cdots \right) dx, \quad -\infty < x < \infty.$$

So we get

$$\int \left(\frac{e^x - 1}{x} \right) dx = C + x + \frac{x^2}{2 \times 2!} + \frac{x^3}{3 \times 3!} + \frac{x^4}{4 \times 4!} + \frac{x^5}{5 \times 5!} + \cdots, \quad -\infty < x < \infty.$$

as required. Note that the radius of convergence is $R = \infty$.

p.95, pr.68

- (b) (14 Points) Find the *radius and interval of convergence* for the series $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}(x+2)^n}{n2^n}$.

Solution: Let $u_n = \frac{(-1)^{n+1}(x+2)^n}{n2^n}$. Then $u_{n+1} = \frac{(-1)^{n+2}(x+2)^{n+1}}{(n+1)2^{n+1}}$ and so

$$\frac{u_{n+1}}{u_n} = \frac{(-1)^{n+2}(-1)^{n+1}(x+2)^{n+1}(x+2)^n}{(n+1)2^{n+1}n2^n} \cdot \frac{n2^n}{(-1)^{n+1}(x+2)^n} = -\frac{1}{2} \frac{n}{n+1} (x+2).$$

Therefore, the power series converges absolutely if

$$\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| < 1 \Rightarrow \lim_{n \rightarrow \infty} \left| -\frac{1}{2} \frac{n}{n+1} (x+2) \right| < 1 \Rightarrow \frac{|x+2|}{2} \lim_{n \rightarrow \infty} \frac{n}{n+1} < 1 \Rightarrow \frac{|x+2|}{2} \underbrace{\lim_{n \rightarrow \infty} \frac{1}{1+1/n}}_1 < 1 \Rightarrow |x+2| < 2,$$

that is, if $-2 < x+2 < 2$, or if, $-4 < x < 0$. Now the endpoints are -4 and 0 . We shall test the series for convergence at these points. When $x = -4$, we have $\sum_{n=1}^{\infty} \frac{-1}{n}$, a divergent series; when $x = 0$, we have $\sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n}$, the alternating harmonic series which converges conditionally. Therefore the radius of convergence is $\boxed{R=2}$ and the interval of convergence is $\boxed{-4 < x \leq 0}$.

p.583, pr.39